CONTENTS

Eduard Hála vi
Vapor–liquid equilibria in ternary mixtures containing nitrogen, methane, ethane and carbon dioxide at low temperatures and high pressures
H. Knapp, X.-D. Yang and Z.-Z. Zhang (Berlin, F.R.G.) 1–18
Simultaneous determination of vapor–liquid equilibria and volumetric properties of ternary systems with a new experimental apparatus
S. Laugier, D. Richon and H. Renon (Paris, France) 19–34
Vapour–liquid equilibrium in the sulphur hexafluoride–n-pentane system at high pressure
J. Berg and Z. Wagner (Prague, Czechoslovakia) 35–45
Vapor–liquid equilibrium measurements of three binary systems made of formic acid, acetic acid and propionic acid by the dew–bubble point temperature method
M. Kato, H. Yoshikawa and M. Yamaguchi (Fukushima, Japan) 47–56
Isobaric vapor–liquid equilibrium in two binary systems involving tert-amyl methyl ether
M. Palczewska-Tulińska, D. Wyrzykowska-Stankiewicz, J. Choliński and K. Zięborak (Warsaw, Poland) 57–68
Vapour–liquid equilibria in the heptane–3-pentanol and heptane–2-methyl-2-butanol systems at constant temperature
J. Wolfová, J. Linek and I. Wichterle (Prague-Suchdol, Czechoslovakia) 69–79
Vapor liquid equilibrium data for systems containing morpholine
C. Pettenati, P. Alessi, M. Fermeglia and I. Kikic (Trieste, Italy) 81–91
Liquid–liquid and vapour–liquid equilibria in the system methyl tert-butyl ether + tetrahydrofuran + water
D. Zikmundová, J. Matouš, J.P. Novák, V. Kubíček and J. Pick (Prague, Czechoslovakia) 93–110
Excess Gibbs energies and excess volumes of the mixtures ethanoic acid + 1,4-dioxane and oxolane
H.E. Kalali, A.M. Demiriz, J. Budde, F. Kohler (Bochum, F.R.G.), A. Dallos and F. Ratkovics (Veszprém, Hungary) 111–120
Infinite-dilution activity coefficients by comparative ebulliometry. Binary systems of chloromethanes, chloroethanes and Freon 113
V. Dohnal and P. Vrbka (Prague, Czechoslovakia) 121–131
Measurement of activity coefficients in highly dilute solutions. Part I
G. Hradetzky, M. Wobst, H. Vopel and H.-J. Bittrich (Merseburg, G.D.R.) 133–145
A comparison of the predictive capability of different group contribution methods
J. Gmehling, D. Tiegs and U. Knipp (Oldenburg, F.R.G.) 147–165
A simplified perturbed hard chain theory equation of state for phase equilibrium calculations for polar fluid mixtures
W. Sheng and B.C.-Y. Lu (Ottawa, Ont., Canada) 167–189

Modification of the extended UNIQUAC model for correlating quaternary liquid–liquid equilibrium data
I. Nagata (Ishikawa, Japan) 191–206
On the standard states for activity coefficients
G. Wilczek-Vera and J.H. Vera (Montreal, Que., Canada) 207–219
Equilibrium behaviour of pure substances from the convex molecule perturbation theory
T. Boublík (Prague, Czechoslovakia) 221–235
Prediction of gas *PVT* data with effective intermolecular potentials using the Haar–Shenker–Kohler equation and computer simulations
B. Saager, A. Lotfi, M. Bohn, V.N. Nguyen and J. Fischer (Bochum. F.R.G.) 237–246
An improved quasilattice expression for liquid phase order–disorder
S. Kemény, Gy. Balog, Gy. Radnai, J. Sawinsky and G. Rezessy (Budapest, Hungary) 247–275
Enthalpies of solution at infinite dilution for linear and branched aliphatic aldehydes in heptane. Calorimetric measurements compared with DISQUAC predictions
M.R. Tiné (Pisa, Italy), G. Della Gatta (Turin, Italy) and H.V. Kehiaian (Paris, France) 277–291
Extension of the group contribution method for the calculation of the heat of vaporization
V. Svoboda and P. Dočkalová (Prague, Czechoslovakia) 293–299
Author index 301